

Local well-posedness of the nonlinear Schrödinger equations on the sphere for data in modulation spaces

Hideo Takaoka*

Abstract

In this paper we discuss a priori estimates derived from the energy method to the initial value problem for the cubic nonlinear Schrödinger on the sphere S^2 . Exploring suitable a priori estimates, we prove the existence of solution for data whose regularity is $s = 1/4$.

1 Introduction

In this paper we consider the initial value problem for the cubic nonlinear Schrödinger equation on the surface:

$$\begin{cases} i\partial_t u + \Delta u = |u|^2, \\ u(x, 0) = u_0(x), \end{cases} \quad (1.1)$$

where the unknown function $u = u(x, t)$ is a complex valued function on $(t, x) \in \mathbb{R} \times S^2$, and Δ denotes the Laplacian on the surface of the unit sphere S^2 in $\mathbb{R}^3$. The equation (1.1) is defocusing. However the result in this paper is independent on whether the equation is focusing or defocussing.

In the case of the whole space case $\mathbb{R}^2$, local well-posedness in $H^s(\mathbb{R}^2)$ for $s \geq 0$ was obtained by Y. Tsutsumi in [16] and T. Cazenave, F. Weissler in [10]. It is important to note that the critical regularity for the scaling argument is $s = 0$. On the other hand, for the periodic boundary problem, local well-posedness for data in $H^s(\mathbb{T}^2)$, $s > 0$ was established by J. Bourgain [2]. In [2], he used the $X^{s,b}$ argument along with the Strichartz estimate which controls the $L^p_{t,x}$ norm of the linear solution in term of the H^s initial data (see also [17]), where this estimate is sharp in the sense that the estimate with $s \leq 0$ fails. For further references, see for instance [4], [15] and references therein.

Among other cases, for the case of two dimensional sphere S^2 , the initial value problem (1.1) is known to be locally well-posed in the Sobolev spaces $H^s(S^2)$

*Department of Mathematics, Hokkaido University, Sapporo, 060-0810, Japan. Email: takaoka@math.sci.hokudai.ac.jp The author was supported in part by J.S.P.S. Grant-in-Aid 25287022.

for $s > 1/4$. This was shown by N. Burq, P. Gérard and N. Tzvetkov [8]¹, where they proved the bilinear Strichartz estimates without loss of derivatives. On the other hand, in [7], they also proved that the flow map is not uniformly continuous in $H^s(S^2)$ for $0 \leq s < 1/4$, although the standard scaling argument concerning the initial value problem (1.1) on the real line case suggests local well-posedness for $s \geq 0$. Further analysis on such instability of solutions was obtained by V. Banica in [11]. It is worth noticing that these instability phenomena arise to be illustrate by considering the following weighted homogeneous spherical harmonic polynomials

$$\psi_k(x_1, x_2, x_3) = k^{1/4-s}(x_1 + ix_2)^k, \quad (x_1, x_2, x_3) \in S^2, \quad k \in \mathbb{N}, \quad (1.2)$$

as initial data. Notice that these weighted homogeneous spherical harmonic polynomials give a concrete counterexample to the restriction estimate obtained by C. D. Sogge [13].

It is natural to ask for the well-posedness result at the regularity $s = 1/4$. In the present paper, we consider some problems at the regularity $s = 1/4$.

We first recall the spherical harmonics on the sphere S^2 .

Notation 1.1. We define $A \lesssim B$ by $A \leq cB$ for some absolute constant $c > 0$. Similarly, we define $A \sim B$ if and only if $A \lesssim B \lesssim A$. We set $\langle a \rangle = (1 + |a|^2)^{1/2}$ for $a \in \mathbb{R}$.

We summarize the properties of eigenvalues and eigenfunctions of the Laplacian Δ on S^2 . Let Y_k^m ($k \in \mathbb{N} \cup \{0\}$, $m \in \mathbb{Z}$, $-k \leq m \leq k$) be complex valued spherical harmonics of degree k and order m . Then Y_k^m are eigenfunctions corresponding to eigenvalues $-\mu_k^2$, where $\mu_k = \sqrt{k(k+1)}$. For a fixed k , we denote by $\mathcal{H}_k$ the $(2k+1)$ -dimensional space spanned by the functions Y_k^m with $-k \leq m \leq k$. We denote by P_k the orthogonal projection on eigenspace $\mathcal{H}_k$.

Let $s \geq 0$. Denote by $H^s = H^s(S^2)$ the Sobolev space associated to $(I - \Delta)^{s/2}$ equipped with norm

$$\|u\|_{H^s} = \left(\sum_{k=0}^{\infty} \langle \mu_k \rangle^{2s} \|P_k u\|_{L^2}^2 \right)^{1/2}.$$

Denote by $B^s = B^s(S^2)$ the Besov type modulation space equipped with norm

$$\|u\|_{B^s} = \sum_{k=0}^{\infty} \langle \mu_k \rangle^s \|P_k u\|_{L^2}.$$

Remark 1.1. (i) We only consider the case when $s = 1/4$, since the argument in this paper will provide the proof in the case $s > 1/4$.

¹In [8], the same problem on a compact Riemann manifold has been discussed.

²The space $L^2(S^2)$ is the Hilbert sum $\bigoplus_{k=0}^{\infty} \mathcal{H}_k$, namely, for every function $f \in L^2(S^2)$, there is a unique Fourier decomposition $f = \sum_{k=0}^{\infty} f_k$ for $f_k \in \mathcal{H}_k$.

(ii) The definition of reasonable Besov space is $B_{p,q}^s$ equipped with norm

$$\|u\|_{B_{p,q}^s} = \left\| 2^{js} \left\| \sum_{2^{j-1} \leq k < 2^j} P_k u \right\|_{L^p(S^2)} \right\|_{\ell_j^q}$$

(see [1]). There is the embedding inclusion relation between B^s and $B_{p,q}^s$, namely

$$B^s \hookrightarrow B_{2,1}^s \hookrightarrow H^s \hookrightarrow B_{2,\infty}^s.$$

From this point of view, the space B^s is a restricted space of the Besov space.

(iii) When $s = 1/4$, the no-weighted homogeneous spherical harmonic functions $\widetilde{\psi}_k(x_1, x_2, x_3) = (x_1 + ix_2)^k$, $(x_1, x_2, x_3) \in S^2$ satisfies $\widetilde{\psi}_k = c_k Y_k^k$ for some constant c_k and

$$P_k \widetilde{\psi}_k = \widetilde{\psi}_k, \quad \widetilde{\psi}_k \in B^{1/4},$$

$$\|\widetilde{\psi}_k\|_{B^{1/4}} = \|\widetilde{\psi}_k\|_{B_{2,1}^{1/4}} = \|\widetilde{\psi}_k\|_{H^{1/4}} = \|\widetilde{\psi}_k\|_{B_{2,\infty}^{1/4}} \sim 1.$$

The main result of this paper is local well-posedness of (1.1) in $B^{1/4}$. More precisely, we show the following theorem.

Theorem 1.1. *For every $r > 0$, there exist $T = T(r) > 0$ and a function space $X_T^{1/4}$ such that given $u_0 \in \{\phi \in B^{1/4} \mid \|\phi\|_{B^{1/4}} < r\}$ there exists a unique solution u of (1.1) with initial data u_0 satisfying $u \in C([0, T]; B^{1/4}) \cap X_T^{1/4}$. Moreover the solution map $\{\phi \in B^{1/4} \mid \|\phi\|_{B^{1/4}} < r\} \ni u_0 \mapsto u \in C([0, T]; B^{1/4}) \cap X_T^{1/4}$ is Lipschitz continuous.*

Remark 1.2. (i) The time of existence T obtained in Theorem 1.1 depends on the value of $\|u_0\|_{B^{1/4}}$. For sequence of initial data $\widetilde{\psi}_k$ ($k \in \mathbb{N}$), Theorem 1.1 guarantees that there exists the sequence of solutions u_k and its existence time T_k . We can say that by Theorem 1.1 the lifespan has the lower estimate $\liminf_{k \rightarrow \infty} T_k \geq c$ for some constant $c > 0$.

(ii) The result in Theorem 1.1 is essentially of the some kind of conditionally local-in-time well-posedness in $B^{1/4}$. Our proof relies on the a priori estimates of solutions. The same sprit in carrying out the proof also works for the unconditionally well-posedness in H^s with $s > 3/4$. However, in this paper we do not discuss at all such a problem.

The well-posedness result for (1.1) is still unclear. But it is interesting to consider the cubic nonlinear Schrödinger equation without gauge invariance. If data are homogeneous spherical harmonic functions, then one can obtain the local well-posedness for data in the Sobolev space H_{hom}^s for $s = 1/4$, where

$$H_{hom}^s = H^s \cap \text{span}\{\psi_k \mid k \geq 0\}.$$

Theorem 1.2. *The initial value problem associated with the cubic nonlinear Schrödinger equation*

$$i\partial_t u + \Delta u = u^3, \quad (1.3)$$

is locally well-posed in $H_{hom}^{1/4}$. Namely, for every $r > 0$, there exist $b > 1/2$, $T = T(r) > 0$ and a function space $X_T^{1/4,b}$ such that given $u_0 \in \{\phi \in H_{hom}^{1/4} \mid \|\phi\|_{H_{hom}^{1/4}} < r\}$ there exists a unique solution u of (1.3) with initial data u_0 satisfying $u \in C([0, T]; H_{hom}^{1/4}) \cap X_T^{1/4,b}$. Moreover the solution map $\{\phi \in H_{hom}^{1/4} \mid \|\phi\|_{H_{hom}^{1/4}} < r\} \ni u_0 \mapsto u \in C([0, T]; H_{hom}^{1/4}) \cap X_T^{1/4,b}$ is Lipschitz continuous.

Remark 1.3. If u_j are smooth in $\text{span}\{\psi_k \mid k \geq 0\}$ for $j = 1, 2, 3$, then the product $\prod_{j=1}^3 u_j$ is within $\text{span}\{\psi_k \mid k \geq 0\}$. This algebraic property allows us to establish the existence of an appropriate class of local solution $u(t) \in C([0, T]; H_{hom}^s)$ of (1.3).

This paper is organized as follows. In section 2, we present the useful function spaces and recall the several estimates for the harmonic projection operators. In section 3, we take the direct estimate for the norm of $\|u(t)\|_{X_T^s}$ along with the energy method under resonance and has several other properties. Multilinear estimates provide us the several type of a priori estimates. In section 4, we carry out the local well-posedness result described in Theorem 1.1. In section 5, we give the proof of Theorem 1.2.

Acknowledgements. I thank Nikolay Tzvetkov for discussion and for pointing out the reference [9].

2 Definition of function spaces and linear estimates

We start out by defining the function spaces.

Definition 2.1. Let $s \geq 0$ and $T > 0$. We define the function space X_T^s as the completion of $C_0^\infty(\mathbb{R} \times S^2)$ under the norm:

$$\|u\|_{X_T^s} = \sum_{k=0}^{\infty} \langle \mu_k \rangle^s \sup_{0 \leq t \leq T} \|P_k u(t)\|_{L^2(S^2)}.$$

We also define the function spaces introduced by Bourgain in [2] and Burq-Gérard-Tzvetkov in [9].

Definition 2.2. For $s \geq 0$ and $b \in \mathbb{R}$, let $X^{s,b}$ be the completion of $C_0^\infty(\mathbb{R}; H^s(S^2))$ for the norm

$$\|f\|_{X^{s,b}} = \left(\sum_{k=0}^{\infty} \|\langle \tau + \mu_k^2 \rangle^b \langle \mu_k \rangle^s \widehat{P_k u}(\tau)\|_{L^2(\mathbb{R} \times S^2)}^2 \right)^{1/2}.$$

Observe that if $b > 1/2$, $X^{s,b} \hookrightarrow C(\mathbb{R}; H^s)$ by usual Sobolev embedding theorem. For $T > 0$, we define the restriction function space

$$X_T^{s,b} = \{f|_{|t| \leq T} \mid f \in X^{s,b}\}.$$

The following extension of the L^p estimates due to Sogge [13] established by Burq, Gérard and Tzvetkov [9] will be a crucial ingredient in our proof.

Theorem 2.1 (Burq, Gérard and Tzvetkov). *There exist $C > 0$ such that, for all $k_1, k_2 \in \mathbb{N} \cup \{0\}$, for all functions f, g on S^2 ,*

$$\|P_{k_1} f P_{k_2} g\|_{L^2(S^2)} \leq C \min\{\langle k_1 \rangle, \langle k_2 \rangle\}^{1/4} \|f\|_{L^2(S^2)} \|g\|_{L^2(S^2)}.$$

Also we shall use the following L^p estimates established by Sogge [13, Theorem 4.2].

Theorem 2.2 (Sogge). *There exist $C > 0$ such that, for all $k \in \mathbb{N} \cup \{0\}$, for all f on S^2 ,*

$$\|P_k u\|_{L^p(S^2)} \leq C \langle k \rangle^{\alpha_p} \|u\|_{L^2(S^2)},$$

where,

- (i) if $2 \leq p \leq 6$, $\alpha_p = (p - 2)/4p$,
- (ii) if $6 \leq p \leq \infty$, $\alpha_p = 2(1/4 - 1/p)$.

3 Nonlinear energy estimates under resonance properties

To show the local well-posedness in $B^{1/4}$, we use the a priori estimates for solution of (1.1). This can be established as follows.

Theorem 3.1. *Let u be a smooth solution of (1.1) on $t \in [0, T]$. Then for all $0 < \delta < 1$, we have*

$$\|u\|_{X_T^{1/4}} \lesssim \|u_0\|_{B^{1/4}} + \left(\sqrt{\frac{T}{\delta}} + \delta^{1/4} \right) \|u\|_{X_T^{1/4}}^2 + \sqrt{T} \|u\|_{X_T^{1/4}}^3.$$

To construct above estimate, first consider a smooth solution u of (1.1) with smooth initial data u_0 . We put $v_k = e^{-it\Delta} P_k u = e^{it\mu_k^2} P_k u$, where we suppressed the t, x dependence and wrote $v_{k_j} = v_{k_j}(t, x)$ for notational simplicity.

Since from the fact that $e^{it\Delta}$ is unitary group in L^2 , we have $\|v_k\|_{L^2} = \|P_k u\|_{L^2}$. Then v_k satisfies the equivalent formula to (1.1):

$$\partial_t v_k = -i P_k \sum_{k_1, k_2, k_3} e^{i\phi(k, k_1, k_2, k_3)t} v_{k_1} \overline{v_{k_2}} v_{k_3}, \quad (3.1)$$

where the phase function $\phi(k, k_1, k_2, k_3)$ is defined by

$$\phi(k, k_1, k_2, k_3) = \mu_k^2 - \mu_{k_1}^2 + \mu_{k_2}^2 - \mu_{k_3}^2.$$

Additionally we may assume that $k \lesssim \max\{k_1, k_2, k_3\}$. Otherwise, the right-hand side of (3.1) is zero, since the $v_{k_1} \overline{v_{k_2}} v_{k_3}$ is the polynomial on degree at most $k_1 + k_2 + k_3$, and it is orthogonal to the polynomial function of degree k .

Then it formally satisfies the identity

$$\begin{aligned} \frac{d}{dt} \|P_k u(t)\|_{L^2}^2 &= 2\operatorname{Re} \int_{S^2} \overline{v_k} \partial_t v_k dx \\ &= 2\operatorname{Im} \int_{S^2} \overline{v_k(t)} \sum_{k_1, k_2, k_3} e^{i\phi(k, k_1, k_2, k_3)t} v_{k_1} \overline{v_{k_2}} v_{k_3} dx. \end{aligned} \quad (3.2)$$

Let us drop the complex number i and simply use c . We decompose (3.2) into a sum of two terms that the resonant contribution $(k_1, k_2, k_3) \in \sigma_k$ and that the non-resonant contribution $(k_1, k_2, k_3) \notin \sigma_k$, where

$$\sigma_k = \left\{ (k_1, k_2, k_3) \in (\mathbb{N} \cup \{0\})^3 : |\mu_k - \sqrt{\mu_{k_1}^2 - \mu_{k_2}^2 + \mu_{k_3}^2}| \leq 1 \text{ or } \mu_k \leq 1/\delta \right\}. \quad (3.3)$$

From the contribution of σ_k , we know that

$$\text{if } |k - k'| > 2 \text{ and } \min\{\mu_k, \mu_{k'}\} > 2/\delta, \text{ then } \sigma_k \cap \sigma_{k'} = \emptyset, \quad (3.4)$$

and

$$\text{if } (k_1, k_2, k_3) \notin \sigma_k, \text{ then } \phi(k, k_1, k_2, k_3) \neq 0. \quad (3.5)$$

Using (3.2), one has after integration in the time interval $[0, t]$ that

$$\begin{aligned} & \|P_k u(t)\|_{B^{1/4}}^2 - \|P_k u_0\|_{B^{1/4}}^2 \\ &= c \int_0^t \int_{S^2} \langle \mu_k \rangle^{\frac{1}{2}} \overline{v_k} \sum_{(k_1, k_2, k_3) \in \sigma_k} e^{i\phi(k, k_1, k_2, k_3)t'} v_{k_1} \overline{v_{k_2}} v_{k_3} dx dt' \\ & \quad + c \int_0^t \int_{S^2} \langle \mu_k \rangle^{\frac{1}{2}} \overline{v_k} \sum_{(k_1, k_2, k_3) \notin \sigma_k} e^{i\phi(k, k_1, k_2, k_3)t'} v_{k_1} \overline{v_{k_2}} v_{k_3} dx dt' \\ &= I_1 + I_2. \end{aligned}$$

Contribution of $(k_1, k_2, k_3) \in \sigma_k$. It is enough to consider the case where k_1 is the maximal modulation of k_j ($j = 1, 2, 3$). By two applications of Theorem 2.1, we obtain

$$\begin{aligned} |I_1| &\lesssim \int_0^t \sum_{(k_1, k_2, k_3) \in \sigma_k} \langle \mu_k \rangle^{\frac{1}{4}} \langle \mu_{k_1} \rangle^{\frac{1}{4}} \|v_k v_{k_2}\|_{L^2} \|v_{k_1} v_{k_3}\|_{L^2} dt' \\ &\lesssim \int_0^t \|v_k\|_{B^{1/4}} \sum_{(k_1, k_2, k_3) \in \sigma_k} \prod_{j=1}^3 \|v_{k_j}\|_{B^{1/4}} dt'. \end{aligned}$$

Contribution of $(k_1, k_2, k_3) \notin \sigma_k$. Since $\phi(k, k_1, k_2, k_3) \neq 0$ by (3.5), we write after integration by parts that

$$\begin{aligned} I_2 &= c \left[\int_{S^2} \frac{\langle \mu_k \rangle^{\frac{1}{2}} e^{i\phi(k, k_1, k_2, k_3)t'}}{\phi(k, k_1, k_2, k_3)} \overline{v_k} \sum_{(k_1, k_2, k_3) \notin \sigma_k} v_{k_1} \overline{v_{k_2}} v_{k_3} \right]_{t'=0}^{t'=t} \\ &\quad + c \int_0^t \int_{S^2} \frac{\langle \mu_k \rangle^{\frac{1}{2}} e^{i\phi(k, k_1, k_2, k_3)t'}}{\phi(k, k_1, k_2, k_3)} \partial_t \left(\overline{v_k} \sum_{(k_1, k_2, k_3) \notin \sigma_k} v_{k_1} \overline{v_{k_2}} v_{k_3} \right) dt' \\ &= I_{21} + I_{22}. \end{aligned}$$

We consider the contribution of I_{21} first. Note that from the non-resonance relation (3.3), we have

$$|\phi(k, k_1, k_2, k_3)| \gtrsim \left\langle \mu_k - \sqrt{\mu_{k_1}^2 - \mu_{k_2}^2 + \mu_{k_3}^2} \right\rangle \langle \mu_k \rangle$$

and $\langle \mu_k \rangle \gtrsim 1/\delta$ within the domain of summation. Then, using Theorem 2.1, we have that the contribution of I_{21} is bounded by

$$\begin{aligned} |I_{21}| &\lesssim \delta^{1-\varepsilon} \|P_k u(t)\|_{B^{1/4}} \sum_{(k_1, k_2, k_3) \notin \sigma_k} \frac{\prod_{j=1}^3 \|P_{k_j} u(t)\|_{B^{1/4}}}{\left\langle \mu_k - \sqrt{\mu_{k_1}^2 - \mu_{k_2}^2 + \mu_{k_3}^2} \right\rangle \langle \mu_k \rangle^\varepsilon} \\ &\quad + \delta^{1-\varepsilon} \|P_k u_0\|_{B^{1/4}} \sum_{(k_1, k_2, k_3) \notin \sigma_k} \frac{\prod_{j=1}^3 \|P_{k_j} u_0\|_{B^{1/4}}}{\left\langle \mu_k - \sqrt{\mu_{k_1}^2 - \mu_{k_2}^2 + \mu_{k_3}^2} \right\rangle \langle \mu_k \rangle^\varepsilon}. \end{aligned}$$

Secondly, we consider the contribution of I_{22} . According to the symmetry, we divide the proof into two cases: the time derivative falls on the factor that v_k and that v_{k_1} .

When the time derivative falls on the factor v_k , $\partial_t v_k$ satisfies the equation (3.1). Apply Theorem 2.1 as above, then the contribution of this case to I_{22} is bounded by

$$\begin{aligned} &c \int_0^t \frac{\|P_k(|u|^2 u)\|_{B^{1/4}} \sum_{(k_1, k_2, k_3) \notin \sigma_k} \prod_{j=1}^3 \|P_{k_j} u\|_{B^{1/4}}}{\left\langle \mu_k - \sqrt{\mu_{k_1}^2 - \mu_{k_2}^2 + \mu_{k_3}^2} \right\rangle \langle \mu_k \rangle} dt' \\ &\lesssim \int_0^t \frac{\|P_k(|u|^2 u)\|_{L^2}}{\langle \mu_k \rangle^{\frac{3}{4}-\varepsilon}} dt' \sup_{0 \leq t' \leq t} \sum_{(k_1, k_2, k_3) \notin \sigma_k} \frac{\prod_{j=1}^3 \|P_{k_j} u\|_{B^{1/4}}}{\left\langle \mu_k - \sqrt{\mu_{k_1}^2 - \mu_{k_2}^2 + \mu_{k_3}^2} \right\rangle \langle \mu_k \rangle^\varepsilon}. \end{aligned}$$

On the other hand, when the time derivative falls on the factor v_{k_1} , we divide two cases that $\mu_k \gtrsim \mu_{k_1}$ and that $\mu_k \ll \mu_{k_1}$.

We consider the contribution of $\mu_k \gtrsim \mu_{k_1}$. By evaluating the weight $\langle \mu_k \rangle^{-1/4}$ by $\langle \mu_{k_1} \rangle^{-1/4}$ in the expression above, the same arguments as in the previous case lead to

$$\left| \int_{S^2} \frac{\overline{P_k u} P_{k_1}(|u|^2 u) \overline{P_{k_2} u} P_{k_3} u}{\langle \mu_k \rangle} dx \right| \lesssim \frac{\|P_k u\|_{B^{1/4}} \|P_{k_1}(|u|^2 u)\|_{L^2} \|P_{k_2} u\|_{B^{1/4}} \|P_{k_3} u\|_{B^{1/4}}}{\langle \mu_k \rangle^\varepsilon \langle \mu_{k_1} \rangle^{\frac{3}{4}-\varepsilon}}.$$

Next we consider the contribution of $\mu_k \ll \mu_{k_1}$. Applying Theorem 2.1 as above, one has that

$$\left| \int_{S^2} \frac{\overline{P_k u} P_{k_1}(|u|^2 u) \overline{P_{k_2} u} P_{k_3} u}{\langle \mu_k - \sqrt{\mu_{k_1}^2 - \mu_{k_2}^2 + \mu_{k_3}^2} \rangle \langle \mu_k \rangle} dx \right| \lesssim \frac{\|P_k u\|_{B^{1/4}} \|P_{k_1}(|u|^2 u)\|_{L^2} \|P_{k_2} u\|_{B^{1/4}} \|P_{k_3} u\|_{B^{1/4}}}{\langle \mu_k - \sqrt{\mu_{k_1}^2 - \mu_{k_2}^2 + \mu_{k_3}^2} \rangle \langle \mu_k \rangle^{\frac{3}{4}}}.$$

We put together all the estimates above. Taking the square root (which is concave) and using Cauchy-Schwarz inequality, we obtain the estimate

$$\begin{aligned} \|u\|_{X_T^{1/4}} &\lesssim \|u_0\|_{B^{1/4}} \\ &+ \left(\sum_{k=0}^{\infty} \|P_k u\|_{L_t([0,T]; B^{1/4})} \right)^{1/2} \left(\int_0^T \sum_{\substack{k \geq 0 \\ (k_1, k_2, k_3) \in \sigma_k}} \prod_{j=1}^3 \|P_{k_j} u(t')\|_{B^{1/4}} dt' \right)^{1/2} \\ &+ \delta^{1/2-\varepsilon} \left(\sum_{k=0}^{\infty} \|P_k u\|_{L_t([0,T]; B^{\frac{1}{4}})} \right)^{\frac{1}{2}} \left(\sum_{\substack{k \geq 0 \\ (k_1, k_2, k_3) \notin \sigma_k}} \frac{\prod_{j=0}^3 \|P_{k_j} u\|_{L_t^\infty([0,T]; B^{1/4})}}{\langle \mu_k - \sqrt{\mu_{k_1}^2 - \mu_{k_2}^2 + \mu_{k_3}^2} \rangle \langle \mu_k \rangle^\varepsilon} \right)^{\frac{1}{2}} \\ &+ \left(\int_0^T \sum_{k=0}^{\infty} \frac{\|P_k(|u|^2 u)\|_{L^2}}{\langle \mu_k \rangle^{\frac{3}{4}-\varepsilon}} dt' \right)^{\frac{1}{2}} \left(\sum_{\substack{k \geq 0 \\ (k_1, k_2, k_3) \notin \sigma_k}} \frac{\prod_{j=1}^3 \|P_{k_j} u\|_{L_t([0,T]; B^{\frac{1}{4}})}}{\langle \mu_k - \sqrt{\mu_{k_1}^2 - \mu_{k_2}^2 + \mu_{k_3}^2} \rangle \langle \mu_k \rangle^\varepsilon} \right)^{\frac{1}{2}} \\ &+ \left(\sum_{k=0}^{\infty} \|P_k u\|_{L_t^\infty([0,T]; B^{\frac{1}{4}})} \right)^{\frac{1}{2}} \times \\ &\quad \times \left(\int_0^T \sum_{\substack{k \gtrsim k_1 \geq 0 \\ (k_1, k_2, k_3) \notin \sigma_k}} \frac{\|P_{k_1}(|u|^2 u)\|_{L^2} \|P_{k_2} u\|_{B^{\frac{1}{4}}} \|P_{k_3} u\|_{B^{\frac{1}{4}}}}{\langle \mu_k - \sqrt{\mu_{k_1}^2 - \mu_{k_2}^2 + \mu_{k_3}^2} \rangle \langle \mu_k \rangle^\varepsilon \langle \mu_{k_1} \rangle^{\frac{3}{4}-\varepsilon}} dt' \right)^{\frac{1}{2}} \\ &+ \left(\sum_{k=0}^{\infty} \|P_k u\|_{L_t^\infty([0,T]; B^{\frac{1}{4}})} \right)^{\frac{1}{2}} \times \\ &\quad \times \left(\int_0^T \sum_{\substack{k_1 \gg k \geq 0 \\ (k_1, k_2, k_3) \notin \sigma_k}} \frac{\|P_{k_1}(|u|^2 u)\|_{L^2} \|P_{k_2} u\|_{B^{\frac{1}{4}}} \|P_{k_3} u\|_{B^{\frac{1}{4}}}}{\langle \mu_k - \sqrt{\mu_{k_1}^2 - \mu_{k_2}^2 + \mu_{k_3}^2} \rangle \langle \mu_k \rangle^{\frac{3}{4}}} dt' \right)^{\frac{1}{2}}, \end{aligned}$$

at least. Since $\sum_{k=0}^{\infty} \|P_k u\|_{L_t([0,T]; B^{1/4})} = \|u\|_{X_T^{1/4}}$, it is sufficient to estimate the following five terms, J_i , $i = 1, \dots, 5$:

$$J_1 = \int_0^T \sum_{\substack{k \geq 0 \\ (k_1, k_2, k_3) \in \sigma_k}} \prod_{j=1}^3 \|P_{k_j} u(t')\|_{B^{1/4}} dt',$$

$$\begin{aligned}
J_2 &= \sum_{\substack{k \geq 0 \\ (k_1, k_2, k_3) \notin \sigma_k}} \frac{\prod_{j=1}^3 \|P_{k_j} u\|_{L_t^\infty([0, T]; B^{1/4})}}{\left\langle \mu_k - \sqrt{\mu_{k_1}^2 - \mu_{k_2}^2 + \mu_{k_3}^2} \right\rangle \langle \mu_k \rangle^\varepsilon}, \\
J_3 &= \int_0^T \sum_{k=0}^\infty \frac{\|P_k(|u|^2 u)\|_{L^2}}{\langle \mu_k \rangle^{\frac{3}{4}-\varepsilon}} dt', \\
J_4 &= \int_0^T \sum_{\substack{k \gtrsim k_1 \geq 0 \\ (k_1, k_2, k_3) \notin \sigma_k}} \frac{\|P_{k_1}(|u|^2 u)\|_{L^2} \|P_{k_2} u\|_{B^{\frac{1}{4}}} \|P_{k_3} u\|_{B^{\frac{1}{4}}}}{\left\langle \mu_k - \sqrt{\mu_{k_1}^2 - \mu_{k_2}^2 + \mu_{k_3}^2} \right\rangle \langle \mu_k \rangle^\varepsilon \langle \mu_{k_1} \rangle^{\frac{3}{4}-\varepsilon}} dt', \\
J_5 &= \int_0^T \sum_{\substack{k_1 \gg k \geq 0 \\ (k_1, k_2, k_3) \notin \sigma_k}} \frac{\|P_{k_1}(|u|^2 u)\|_{L^2} \|P_{k_2} u\|_{B^{\frac{1}{4}}} \|P_{k_3} u\|_{B^{\frac{1}{4}}}}{\left\langle \mu_k - \sqrt{\mu_{k_1}^2 - \mu_{k_2}^2 + \mu_{k_3}^2} \right\rangle \langle \mu_k \rangle^{\frac{3}{4}}} dt'.
\end{aligned}$$

Subsequently, we estimate J_i for each separately.

Estimate of J_1 . For $\mu_k > 1/\delta$, the each support σ_k is disjoint by (3.4). Therefore J_2 is bounded by

$$J_1 \lesssim \frac{T}{\delta} \sum_{k_1, k_2, k_3} \prod_{j=1}^3 \|P_{k_j} u\|_{L_t^\infty([0, T]; B^{\frac{1}{4}})} \lesssim \frac{T}{\delta} \|u\|_{X_T^{\frac{1}{4}}}^3.$$

Estimate of J_2 . Due to $\left\langle \mu_k - \sqrt{\mu_{k_1}^2 - \mu_{k_2}^2 + \mu_{k_3}^2} \right\rangle^{-1} \langle \mu_k \rangle^{-\varepsilon} \in \ell_k^1$, we have that J_2 is bounded by

$$J_2 \lesssim \sum_{k_1, k_2, k_3} \prod_{j=1}^3 \|P_{k_j} u\|_{L_t^\infty([0, T]; B^{\frac{1}{4}})} \lesssim \|u\|_{X_T^{\frac{1}{4}}}^3.$$

Estimate of J_3 . Cauchy-Schwarz inequality provides $J_3 \lesssim \|u\|_{L_t([0, T], L_x^6)}^3$. Then we have that

$$J_3 \lesssim T \left\| \sum_{k=0}^\infty P_k u \right\|_{L_t^\infty([0, T]; L^6)}^3 \lesssim \left(\sum_{k=0}^\infty \|P_k u\|_{L_t^\infty([0, T]; L^6)} \right)^3.$$

Now we use Sobolev inequality in Theorem 2.2 to obtain

$$J_3 \lesssim T \left(\sum_{k=0}^\infty \langle \mu_k \rangle^{\frac{1}{6}} \|P_k u\|_{L_t^\infty([0, T]; L^2)} \right)^3 \lesssim T \|u\|_{X_T^{\frac{1}{6}}}^3.$$

Estimate of J_4 . It suffices to show that

$$\sup_{0 \leq t \leq T} \sum_{\substack{k \gtrsim k_1 \geq 0 \\ (k_1, k_2, k_3) \notin \sigma_k}} \frac{\|P_{k_1}(|u|^2 u)\|_{L^2} \|P_{k_2} u\|_{B^{\frac{1}{4}}} \|P_{k_3} u\|_{B^{\frac{1}{4}}}}{\left\langle \mu_k - \sqrt{\mu_{k_1}^2 - \mu_{k_2}^2 + \mu_{k_3}^2} \right\rangle \langle \mu_k \rangle^\varepsilon \langle \mu_{k_1} \rangle^{\frac{3}{4}-\varepsilon}} \lesssim \|u\|_{X_T^{\frac{1}{4}}}^5.$$

The sum Σ_k is uniformly bounded due to $\left\langle \mu_k - \sqrt{\mu_{k_1}^2 - \mu_{k_2}^2 + \mu_{k_3}^2} \right\rangle^{-1} \langle \mu_k \rangle^{-\varepsilon} \in \ell_k^1$. Now we apply same argument as in the previous proof for J_3 . Therefore

$$J_4 \lesssim T \|u\|_{X_T^{\frac{1}{4}}}^5.$$

Estimate of J_5 . A simple calculation shows that J_5 is bounded by

$$J_5 \lesssim T \sup_{0 \leq t \leq T} \sum_{k_1, k_2, k_3} \frac{\|P_{k_1}(|u|^2 u)\|_{L^2} \|P_{k_2} u\|_{B^{\frac{1}{4}}} \|P_{k_3} u\|_{B^{\frac{1}{4}}}}{\langle \mu_{k_1}^2 - \mu_{k_2}^2 + \mu_{k_3}^2 \rangle^{\frac{3}{8} - \varepsilon}}.$$

Now by Cauchy-Schwarz inequality with the fact $\langle \mu_{k_1}^2 + \mu_{k_2}^2 + \mu_{k_3}^2 \rangle^{-3/8 + \varepsilon} \in \ell_{k_1}^2$, we see that

$$J_5 \lesssim T \|u\|_{L_t([0, T]; L^6)}^3.$$

From the previous step in the proof for J_3 , we have that

$$J_5 \lesssim T \|u\|_{X_T^{\frac{1}{4}}}^5.$$

This completes the proof of Theorem 3.1. $\square$

Repeating a similar computation on the difference of two solutions, we have the following theorem.

Theorem 3.2. *Let u and v be two smooth solutions of (1.1) on $t \in [0, T]$ with initial data u_0 and v_0 , respectively. Then for all $0 < \delta < 1$, we have*

$$\|u - v\|_{X_T^{\frac{1}{4}}} \lesssim \|u_0 - v_0\|_{B^{\frac{1}{4}}} + \left(\sqrt{\frac{T}{\delta}} + \delta^{1/4} \right) M \|u - v\|_{X_T^{\frac{1}{4}}} + \sqrt{T} M^2 \|u - v\|_{X_T^{\frac{1}{4}}}.$$

where $M = \max \left\{ \|u\|_{X_T^{1/4}}, \|v\|_{X_T^{1/4}} \right\}$.

4 Proof of Theorem 1.1

Now we explain how to use Theorems 3.1 and 3.2 to establish local solutions.

Let $u_{0,n}$ be a sequence in $H^\infty(S^2)$ such that $u_{0,n} \rightarrow u_0$ in $B^{1/4}$ as $n \rightarrow \infty$. For $u(x, 0) = u_{0,n}(x)$ in (1.1), we have global smooth solutions of (1.1) which we denote by u_n .

Let $T > 0$ be a positive constant with $T < 1$ to be determined later. First we use the continuity argument to show that u_n satisfies the a priori estimate obtained in Theorem 3.1 on the time interval $[0, T]$ independent of n . Namely, we have

$$\|u_n\|_{X_T^{1/4}} \leq C \|u_{0,n}\|_{B^{1/4}} + C \left(\sqrt{\frac{T}{\delta}} + \delta^{1/4} \right) \|u_n\|_{X_T^{1/4}}^2 + C \sqrt{T} \|u_n\|_{X_T^{1/4}}^3.$$

We choose $R > 0$ so large that $\|u_0\|_{B^{1/4}} \leq R$, $\|u_{0,n}\|_{B^{1/4}} \leq R$ for all n . Since $\|u_n\|_{X_t^{1/4}}$ is continuous in t , there exist small $\delta > 0$ and a time interval $[0, T']$ such that $\|u_n\|_{X_{T'}^{1/4}} \leq 2CR$. Then we have by Theorem 3.1

$$\begin{aligned} \|u_n\|_{X_{T'}^{1/4}} &\leq C\|u_{0,n}\|_{B^{1/4}} + C\left(\sqrt{\frac{T'}{\delta}} + \delta^{1/4}\right)\|u_n\|_{X_{T'}^{1/4}}^2 + C\sqrt{T'}\|u_n\|_{X_{T'}^{1/4}}^3 \\ &\leq CR + \frac{CR}{2} = \frac{3}{2}CR. \end{aligned}$$

Therefore we can choose $T < 1$ so small that

$$\|u_n\|_{X_T^{1/4}} \leq 2R$$

where T depends only on C and R .

Moreover, we repeat a similar argument on difference of two solutions. Theorem 3.2 yields

$$\|u_n - u_m\|_{X_T^{1/4}} \leq C'\|u_{0,n} - u_{0,m}\|_{B^{1/4}}$$

for some $C' > 0$. Hence by the Ascoli-Arzel'a compactness theorem we construct convergent sequences $\{u_{n(k)}\} \subset \{u_n\}$ by taking subsequences. Thus we obtain a solution u of (1.1) satisfying

$$u \in X_T^{1/4} \subset L^\infty([0, T]; B^{1/4}), \quad \|u\|_{X_T^{1/4}} \leq 2R,$$

$$\lim_{k \rightarrow \infty} \|u_{n(k)} - u\|_{X_T^{1/4}} = 0.$$

We now present the proof of uniqueness of solution to (1.1). Let u and v be two solutions of (1.1) in $X_T^{1/4}$ with the same initial data u_0 . We evaluate the $X_T^{1/4}$ norm of $u - v$ in the same way as above by using Theorem 3.2. If we choose $\delta > 0$, $T' > 0$ small, then

$$\|u - v\|_{X_{T'}^{1/4}} \leq \eta\|u - v\|_{X_{T'}^{1/4}}$$

for some $0 < \eta < 1$, which shows $u = v$ on $[0, T']$. By repeating this procedure, we obtain $u = v$ on $[0, T]$.

Moreover repeating the proof of Theorem 3.1 with the functional

$$\|P_k u(t_1) - P_k u(t_2)\|_{L^2}^2 = \int_{t_2}^{t_1} \frac{d}{dt'} \|P_k u(t') - P_k u(t_2)\|_{L^2}^2 dt',$$

we obtain that

$$\|u(t_2) - u(t_1)\|_{B^{1/4}} \leq C\left(\sqrt{\frac{t_1 - t_2}{\delta}} + \delta^{1/4}\right)\|u\|_{X_T^{1/4}}^2 + C\sqrt{t_1 - t_2}\|u\|_{X_T^{1/4}}^3$$

whenever $0 \leq t_2 \leq t_1 \leq T$ and $0 < \delta < 1$. Thus if $t_1 \rightarrow t_2$ and $\delta \rightarrow 0$, that $u(t_2) \rightarrow u(t_1)$ in $B^{1/4}$. Hence $u \in C([0, T]; B^{1/4})$.

The continuous dependence of solution on initial data is proven in the same way as in the proof of existence of solution. $\square$

5 Proof of Theorem 1.2

In this section, we will solve the integral equation

$$u(t) = e^{it\Delta}u_0 - i \int_0^t e^{i(t-t')\Delta}u^3(t') dt', \quad u_0 \in H_{hom}^{1/4},$$

in $C([0, T]; H_{hom}^{1/4}) \cap X_T^{1/4, b}$ for appropriate $b > 1/2$.

Remark 5.1. A similar argument as in [9] provides us the local well-posedness result of the Cauchy problem (1.3) with no change to their proof. More precisely, if $u_0 \in H^s$ with $s > 1/4$, there exists a unique solution $u(t) \in H^s$ for local in time. As it was remarked in Remark 1.3, it is easy to see that we can have the local result in the homogeneous space H_{hom}^s for $s > 1/4$.

We summarize the properties of the space $X^{s, b}$.

Lemma 5.1. *Let $s \geq 0$, $1/2 < b < b' < 1$ and $0 < T < 1$. There exists $c > 0$ such that*

$$\|\theta_T(t)e^{it\Delta}u_0\|_{X^{s, b}} \leq cT^{1/2-b}\|u_0\|_{H^s},$$

$$\left\| \theta(t) \int_0^t e^{i(t-t')\Delta} f(t') dt' \right\|_{X^{s, b}} \leq c\|f\|_{X^{s, b-1}},$$

$$\|\theta_T f\|_{X^{s, b}} \leq cT^{1/2-b}\|f\|_{X^{s, b}},$$

and

$$\|\theta_T f\|_{X^{s, b-1}} \leq cT^{b'-b}\|f\|_{X^{s, b'-1}},$$

where $\theta \in C_0^\infty(\mathbb{R})$, $0 \leq \theta \leq 1$, $\theta = 1$ near 0, $\text{supp}\theta \subset [-1, 1]$, and $\theta_T(t) = \theta(T^{-1}t)$.

The proof of Lemma 5.1 is given by Bourgain in [2], and Burq-Gérard-Tzvetkov in [9].

We need to have an estimate that takes the nonlinearity u^3 in $X^{s, b-1}$. We comment on the bilinear Strichartz estimate constructed in [9], and prove Theorem 1.2.

Lemma 5.2. *Let $b > 3/8$ and $f_j \in X^{1/4, b}$ be homogeneous spherical functions of the form*

$$f_j(t, x) = \sum_{k \geq 0} a_{j, k}(t) \sin^k \theta e^{ik\phi}, \quad (5.1)$$

for $j = 1, 2$, where θ, ϕ are spherical coordinates $x = (\sin \theta \cos \phi, \sin \theta \sin \phi, \cos \theta)$, $0 \leq \theta < \pi$, $0 \leq \phi < 2\pi$. Then the following bilinear estimate holds:

$$\|f_1 f_2\|_{L_{t, x}^2} \lesssim \|f_1\|_{X^{0, b}} \|f_2\|_{X^{1/4, b}}.$$

For the proof of Lemma 5.2, we will need some elementary inequalities.

Lemma 5.3. *Let $0 < \alpha \leq \beta$ such that $\alpha + \beta > 1$ and $\varepsilon > 0$. Then*

$$\int_{-\infty}^{\infty} \frac{dt}{\langle t-a \rangle^{\alpha} \langle t-\beta \rangle^{\beta}} \lesssim \frac{1}{\langle \alpha-\beta \rangle^{\gamma}},$$

where

$$\gamma = \begin{cases} \alpha + \beta - 1, & \text{if } \beta < 1, \\ \alpha - \varepsilon, & \text{if } \beta = 1, \\ \alpha, & \text{if } \beta > 1. \end{cases}$$

For the proof of Lemma 5.3, see [12] for instance.

Proof of Lemma 5.2. The proof follows the same general outline as arguments by Bourgain [2] and C. E. Kenig, G. Ponce and L. Vega [12].

By the Parseval identity in time variable, we have

$$\|f_1 f_2\|_{L^2_{t,x}} = \left\| \int_{\mathbb{R}} \sum_{k_1, k_2 \geq 0} \widehat{a_{1,k_1}}(\tau - \tau_1) \widehat{a_{2,k_2}}(\tau_1) \sin^{k_1+k_2+1/2} \theta e^{i(k_1+k_2)\phi} d\tau_1 \right\|_{L^2_{\tau,\theta,\phi}}.$$

Using the similar computation to [2], we have that $\|f_1 f_2\|_{L^2}^2$ is equal to

$$\begin{aligned} & \sum_{k_1, k'_1, k_2, k'_2} \int_{\mathbb{R}^3} \int_0^{2\pi} \int_0^{\pi} \widehat{a_{1,k_1}}(\tau - \tau_1) \widehat{a_{2,k_2}}(\tau_1) \overline{\widehat{a_{1,k'_1}}(\tau - \tau'_1) \widehat{a_{2,k'_2}}(\tau'_1)} \\ & \times \sin^{k_1+k'_1+k_2+k'_2+1} \theta e^{i(k_1+k_2-k'_1-k'_2)\phi} d\tau_1 d\tau'_1 d\tau d\theta d\phi. \end{aligned}$$

It is easy to see that

$$\int_0^{2\pi} e^{i(k_1+k_2-k'_1-k'_2)\phi} d\phi = 2\pi \delta(k_1+k_2-k'_1-k'_2),$$

and

$$\int_0^{\pi} \sin^{k_1+k'_1+k_2+k'_2+1} \theta d\theta \sim \frac{1}{\sqrt{k_1+k'_1+k_2+k'_2+1}}.$$

We write $m = k_1 + k_2 = k'_1 + k'_2$ and so we can conclude that

$$\|f_1 f_2\|_{L^2}^2 \lesssim \sum_m \frac{1}{\sqrt{2m+1}} \int_{\mathbb{R}} \left(\sum_{k_1} \int_{\mathbb{R}} |\widehat{a_{1,k_1}}(\tau - \tau_1) \widehat{a_{2,m-k_1}}(\tau_1)| d\tau_1 \right)^2 d\tau.$$

Using Cauchy-Schwarz, the inner integral can be estimated as follows:

$$\begin{aligned} & \sum_{k_1} \int_{\mathbb{R}} |\widehat{a_{1,k_1}}(\tau - \tau_1) \widehat{a_{2,m-k_1}}(\tau_1)| d\tau_1 \\ & \lesssim M \left(\sum_{k_1} \int_{\mathbb{R}} |\widehat{a_{1,k_1}}(\tau - \tau_1)|^2 \langle \tau - \tau_1 - \mu_{k_1}^2 \rangle^{2b} |\widehat{a_{2,m-k_1}}(\tau_1)|^2 \langle \tau_1 - \mu_{m-k_1}^2 \rangle^{2b} d\tau_1 \right)^{1/2}, \end{aligned}$$

where

$$M = \sup_{\tau \in \mathbb{R}, m \geq 0} \left(\sum_{k_1} \int_{\mathbb{R}} \langle \tau - \tau_1 - \mu_{k_1}^2 \rangle^{-2b} \langle \tau_1 - \mu_{m-k_1}^2 \rangle^{-2b} d\tau_1 \right)^{1/2}.$$

A simple computation by Lemma 5.3 yields

$$M^2 \lesssim \sup_{\tau, m} \sum_{k_1} \langle \tau - \mu_{k_1}^2 - \mu_{m-k_1}^2 \rangle^{-4b+1}.$$

For each τ and m , we obtain the two roots $k_1 = \alpha_{\pm}(\tau, m)$ of the quadratic equation with respect to k_1 :

$$\tau - \mu_{k_1}^2 - \mu_{m-k_1}^2 = 0.$$

Then we decompose the sum of $\sum_{k_1}$ into two cases:

- $|k_1 - \alpha_+(\tau, m)| < 1$ or $|k_1 - \alpha_-(\tau, m)| < 1$,
- $|k_1 - \alpha_{\pm}(\tau, m)| \geq 1$.

Therefore we conclude that

$$M^2 \lesssim 1 + \sup_{\tau, m} \sum_{|k_1 - \alpha_{\pm}(\tau, m)| \geq 1} \langle k_1 - \alpha_+(\tau, m) \rangle^{-4b+1} \langle k_1 - \alpha_-(\tau, m) \rangle^{-4b+1} \lesssim 1,$$

since by $b > 3/8$.

Then it follows that $\|f_1 f_2\|_{L^2}^2$ is bounded by

$$\begin{aligned} & c \sum_{m, k_1} \frac{1}{\sqrt{2m+1}} \int_{\mathbb{R}^2} |\widehat{a_{1, k_1}}(\tau - \tau_1)|^2 \langle \tau - \tau_1 - \mu_{k_1}^2 \rangle^{2b} |\widehat{a_{2, m-k_1}}(\tau_1)|^2 \langle \tau_1 - \mu_{m-k_1}^2 \rangle^{2b} d\tau_1 d\tau \\ & \lesssim \sum_{k_1} \int_{\mathbb{R}} \frac{|\widehat{a_{1, k_1}}(\tau_1) \langle \tau_1 - \mu_{k_1}^2 \rangle^b|^2}{\langle \mu_{k_1} \rangle^{1/4}} d\tau_1 \sum_{k_2} \int_{\mathbb{R}} |\widehat{a_{2, k_2}}(\tau_2) \langle \tau_2 - \mu_{k_2}^2 \rangle^b|^2 d\tau_2 \\ & = c \|f_1\|_{X^{0, b}}^2 \|f_2\|_{X^{1/4, b}}^2, \end{aligned}$$

which completes the proof. $\square$

Lemma 5.2 is equivalent to proving the following result.

Lemma 5.4. *Let $f_j \in X^{1/4, 1/2+}$ ($1 \leq j \leq 3$) be of form in (5.1). Then for all $b' < 5/8$, $b > 3/8$, the trilinear estimate*

$$\left\| \prod_{j=1}^3 f_j \right\|_{X^{1/4, b'-1}} \lesssim \prod_{j=1}^3 \|f_j\|_{X^{1/4, b}} \quad (5.2)$$

holds.

Proof. First notice that by Remark 1.3 the product function $\prod_{j=1}^3 f_j$ preserves of form in (5.1). Indeed, one has that there exist $a_k(t)$ ($k \geq 0$) such that

$$\prod_{j=1}^3 f_j(t, x) = \sum_{k \geq 0} a_k(t) \sin^k \theta e^{ik\phi}.$$

Using duality technique, orthogonality argument, Leibniz rule for fractional derivatives and symmetry argument, it suffices to show that

$$\left| \int_{\mathbb{R} \times S^2} f_4(t, x) \prod_{j=1}^3 f_j(t, x) dt dx \right| \lesssim \|f_1\|_{X^{1/4, b}} \|f_2\|_{X^{1/4, b}} \|f_3\|_{X^{0, b}} \|f_4\|_{X^{0, 1-b'}} \quad (5.3)$$

for any $f_4 \in X^{0, 1-b'}$ of form

$$f_4(t, x) = \sum_{k \geq 0} a_{4, k}(t) \sin^k \theta e^{ik\phi}.$$

By two applications of Lemma 5.2, we have that the left-hand side of (5.3) is bounded by

$$\|f_1 f_3\|_{L_{t, x}^2} \|f_2 f_4\|_{L_{t, x}^2} \lesssim \|f_1\|_{X^{1/4, b}} \|f_3\|_{X^{0, b}} \|f_2\|_{X^{1/4, b}} \|f_4\|_{X^{0, 1-b'}},$$

provided $b > 3/8$ and $b' < 5/8$. Thus lemma follows. $\square$

Proof of Theorem 1.2. For positive constants T and r , we define

$$E(T, r) = \left\{ u \in C_0(\mathbb{R}; H_{hom}^{1/4}) \cap X^{s, b} \mid \|\theta_T u\|_{X^{1/4, b}} \leq r T^{1/2-b} \right\}.$$

We shall show that for appropriate values of $T > 0$ and $b > 1/2$, the following map $u \rightarrow \Phi(u)$ defines a contraction map on $E(T, r)$ for appropriate values of T and r :

$$\Phi(u)(t) = e^{it\Delta} u_0 - i \int_0^t e^{i(t-t')\Delta} [\theta_T(\theta_T u^3)](t') dt'. \quad (5.4)$$

Using Lemmas 5.1 and 5.4 combined with the integral formula in (5.4), we obtain

$$\begin{aligned} \|\theta_T \Phi(u)\|_{X^{1/4, b}} &\leq c T^{1/2-b} \|u\|_{H^{1/4}} + c T^{1/2-b} \|\theta_T(\theta_T u)^3\|_{X^{1/4, b-1}} \\ &\leq c T^{1/2-b} \|u_0\|_{H^{1/4}} + c T^{b'-b+1/2-b} \|(\theta_T u)^3\|_{X^{1/4, b'-1}} \\ &\leq c T^{1/2-b} \|u_0\|_{H^{1/4}} + c T^{b'-b+1/2-b} \|\theta_T u\|_{X^{1/4, b}}^3 \\ &\leq c T^{1/2-b} \|u_0\|_{H^{1/4}} + c T^{b'-b+3(1/2-b)} r^3 \end{aligned}$$

for $1/2 < b < b' < 5/8$. Set $r = 2c\|u_0\|_{H^{1/4}}$. Choosing $T > 0$ small and $b' - b + 3(1/2 - b) > 0$ such that $T^{b'-b+2(1/2-b)} r^2 \leq 1/2$, we have that

$$\|\theta_T \Phi(u)\|_{X^{1/4, b}} \leq r T^{1/2-b}.$$

A similar argument to above shows that

$$\|\theta_T \Phi(u_1) - \theta_T \Phi(u_2)\|_{X^{1/4,b}} \leq \frac{1}{2} \|\theta_T u_1 - \theta_T u_2\|_{X^{1/4,b}}.$$

Notice that $X^{s,b} \hookrightarrow C(\mathbb{R}; H^s)$ when $b > 1/2$. Therefore, a standard iteration argument shows that we obtain a unique solution u of (1.3) satisfying

$$\theta_T u \in ([0, T]; H_{hom}^s) \cap X^{1/4,b}, \quad \|\theta_T u\|_{X^{1/4,b}} \leq rT^{1/2-b}.$$

This proves Theorem 1.2 □

References

- [1] J. Bergh and J. Löfström, *Interpolation spaces: an introduction*, Springer, 1966.
- [2] J. Bourgain, *Fourier transform restriction phenomena for certain lattice subsets and applications to nonlinear evolution equations, I, II*, Geom. Funct. Anal., **3**, 1993, 107–156, 209–262.
- [3] J. Bourgain, *Refinements of Strichartz’ inequality and applications to 2D-NLS with critical nonlinearity*, International Mathematics Research Notices, **5**, 1998, 253–283.
- [4] J. Bourgain, *Global solutions of nonlinear Schrödinger equations*, Amer. Mathematical Society, 1999.
- [5] J. Bourgain and A. Bulut, *Almost sure global well posedness for the radial nonlinear Schrödinger equation on the unit ball I: The 2D case*, Ann. Inst. H. Poincaré Anal. Non Linéaire, **31**, 2014, 1267–1288.
- [6] J. Bourgain and A. Bulut, *Almost sure global well-posedness for the radial nonlinear Schrödinger equation on the unit ball II: the 3d case*, J. Eur. Math. Soc., **16**, 2014, 1289–1325.
- [7] N. Burq, P. Gerard and N. Tzvetkov, *An instability property of the nonlinear Schrödinger equation on S^d* , Math. Res. Lett., **9**, 2002, 323–335.
- [8] N. Burq, P. Gerard and N. Tzvetkov, *Strichartz inequalities and the nonlinear Schrödinger equation on compact manifolds*, Amer. J. Math., **126**, 2004, 569–605.
- [9] N. Burq, P. Gerard and N. Tzvetkov, *Bilinear eigenfunction estimates and the nonlinear Schrödinger equation on surfaces*, Invent. Math., **159**, 2005, 187–223.
- [10] T. Cazenave and F. Weissler, *The Cauchy problem for the critical nonlinear Schrödinger equation in H^s* , Nonlinear Analysis, Theory, Methods and Applications, **14**, 1990, 807–836.

- [11] Z. Hani, *Global well-posedness of the cubic nonlinear Schrödinger equation on closed manifolds*, Commun. in Partial Diff. Equations, **37**, 2012, 1186–1236.
- [12] C. E. Kenig, G. Ponce and L. Vega, *A bilinear estimate with applications to the KdV equation*, J. Amer. Math. Soc., **9**, 1996, 573–603.
- [13] C. D. Sogge, *Oscillatory integrals and spherical harmonics*, Duke Math. J., **53**, 1986, 43–65.
- [14] C. D. Sogge, *Fourier integrals in classical analysis*, Cambridge University Press, 1993.
- [15] T. Tao, *Nonlinear dispersive equations: local and global analysis*, Amer. Mathematical Society, 2006.
- [16] Y. Tsutsumi, *L^2 solutions for nonlinear Schrödinger equations and nonlinear groups*, Funk. Ekva., **30**, 1987, 115–125.
- [17] A. Zygmund, *On Fourier coefficients and transforms of functions of two variables*, Studia Math., **50**, 1974, 189–201.